



Continuing Education Questionnaire

Topic: Mood Disorders in TBI

Date: April 29, 2022

Write Your Name: _____

Once this form is completed, please submit to institute@sevitahealth.com

1. What percentage of patients who have sustained a traumatic brain injury patients will develop mood disorders?

- a. 10%
- b. 99%
- c. 50%
- d. 20%

2. What is the gold standard for diagnosing post traumatic mood disorders?

- a. DSM
- b. PCL
- c. PHQ-9
- d. GAD-7

3. What is a good screening tool for anxiety?

- a. PHQ-9
- b. Western aphasia battery
- c. ABS
- d. GAD-7



Continuing Education Questionnaire

4. The ____ are considered the first line pharmacologic treatment options for post traumatic depression?
- a. SSRIs
 - b. SNRIs
 - c. Anxiolytics
 - d. Benzodiazepines
5. The Agitated Behavior scale (ABS) is used to track what?
- a. Post traumatic insomnia
 - b. Bladder incontinence
 - c. Post traumatic agitation
 - d. Post traumatic headaches
6. Pseudobulbar affect (PBA) pathological laughter and ____.
- a. Sweating
 - b. Tachycardia
 - c. Crying
 - d. Verbal outbursts
7. What is one of the two most common lobes of the brain that are injured in head trauma??
- a. Frontal lobe
 - b. Occipital lobe
 - c. Parietal lobe
 - d. Pineal gland
8. What is a screening tool for PTSD?
- a. ABS
 - b. PHQ-9
 - c. Parietal lobe
 - d. Pineal gland



Continuing Education Questionnaire

9. The young mania rating scale can be used for what mood disorder?

- a. Depression
- b. Bipolar disorder
- c. Anxiety
- d. Agitation

10. Premorbid alcohol abuse is considered a risk factor for ____ after sustaining a TBI.

- a. Crying
- b. Depression
- c. Tachycardia
- d. Hypoxia